



Discover Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

Celebrating 35 Years of the AJNR: March 1987 edition

AJNR Am J Neuroradiol 2022, 43 (3) 507

doi: <https://doi.org/10.3174/ajnr.P6841>

<http://www.ajnr.org/content/43/3/507.citation>

This information is current as
of June 14, 2025.

Celebrating 35 Years of the AJNR

March 1987 edition

Internuclear Ophthalmoplegia: MR-Anatomic Correlation

Scott W. Atlas¹
Robert L. Grossman¹
Peter J. Savino²
Norman J. Schatz²
Robert C. Sergott³
Thomas M. Bosley⁴
David B. Hackney¹
Herbert I. Goldberg¹
Larissa T. Blaniuk¹
Robert A. Zimmerman¹

Internuclear ophthalmoplegia is a gaze disorder characterized by impairment on the side of a lesion involving the medial longitudinal fasciculus with nystagmus of the abducting eye. Eleven patients with internuclear ophthalmoplegia (nine with clinical multiple sclerosis, two with clinical infarction) underwent with spin-echo techniques on a 1.5-T system. Nine patients also had CT. focal or nodular areas of high signal intensity on T2-weighted images in the medial longitudinal fasciculus in 10 of 11 patients. In one of four patients with internuclear ophthalmoplegia who had MRI after intravenous gadolinium-DTPA, a ring lesion was seen in the region of the medial longitudinal fasciculus on T1-weighted images, indicating active blood-brain-barrier disruption, which correlates with this patient's recent-onset internuclear ophthalmoplegia. CT failed to show in all nine patients examined. This report demonstrates the superiority of MRI in detecting gaze disorders attributable to brainstem dysfunction, such as internuclear ophthalmoplegia, and correlates MR findings with the relevant neuroanatomy of longitudinal fasciculus.

Internuclear ophthalmoplegia (INO) is a disorder of eye movement, characterized by impaired adduction on the side of a lesion involving longitudinal fasciculus (MLF) with dissociated nystagmus of the abducting eye [1, 2]. In young patients, this syndrome is most commonly caused by multiple sclerosis (MS) [1, 3-6]; in fact, INO is the most common oculomotor manifestation of MS [3]. The same eye-movement disorder in an older age group is usually attributed to cerebrovascular disease. MR is a highly sensitive imaging technique for detecting MS plaques [7-10], and it has become the technique of choice for



The Role of Dural Anomalies in Vein of Galen Aneurysms: Report of Six Cases and Review of the Literature

P. Lasjaunias^{1,2}
K. Ter Brugge³
L. Lopez-Ibor⁴
M. Chiu²
O. Fiodmark⁵
S. Chuang⁶
J. Gossage⁷

It is proposed that the vein of Galen aneurysm is an increased flow (usually caused by a direct or indirect connection of the vein of Galen aneurysm associated with obstruction of a dural sinus venous obstruction is to the vein of Galen, obstructive (noncommunicating) hydrocephalus drainage from the rest of the brain will be unaltered from the vein of Galen aneurysm, the communicating type of hydrocephalus. The magnitude of the arteriovenous shunt. It may be related to the retrograde venous flow in the healthy brain. Careful attention should be paid of the lesion because the types of dural venous case to case. The term "vein of Galen aneurysm" term "vein of Galen ectasia."

Since concluding our research in infants, children [1], we have been involved in the study of vein of Galen aneurysm (VGA). We report six cases of associated dural anomalies. Careful analysis of deep-seated arteriovenous malformations (AVMs) and their relationship to the VGA is discussed.

The purpose of the present paper is to contribute to the angioanatomic analysis of the VGA, to discuss what may be the primary developmental defect, and to consider the relationship between the anomalies encountered and the clinical outcome of the disease.




Case Reports

Case 1

A 4-kg newborn girl presented with cardiac failure. Vertebral angiography demonstrated a VGA with agenesis of the straight sinus and high-flow venous collateral circulation from the VGA to the internal carotid (Fig. 1). There was angiographic evidence of arterial steal, but there was no retrograde venous drainage into the cortical veins or sinuses.

Case 2

A 7-kg infant boy presented with an increase in head circumference. A vertebral angiogram showed a VGA with agenesis of the straight sinus and moderately high-flow collateral circulation from the VGA to the lateral and cavernous sinuses (Fig. 2). Angiographically, no arterial steal was demonstrated, but retrograde venous drainage into the cortical veins was noted.



Received June 21, 1986; accepted September 21, 1986.
¹ Department of Radiology, University of Pennsylvania, Philadelphia, PA 19104, Grossman.
² Neurophysiology, University of Pennsylvania, Philadelphia, PA 19104, Grossman.
AJNR 8:243-247, 1987.
© American Society of Neuroradiology.

Received March 3, 1986; accepted after revision September 11, 1986.

Presented at the 24th annual meeting of the American Society of Neuroradiology, San Diego, January 1986.

¹ Service de Radiologie, Hôpital de Bicêtre, 78, Rue du Général Leclerc, 94270 Le Kremlin-Bicêtre, France. Address reprint requests to P. Lasjaunias.

² Department of Radiology, Toronto Western Hospital, Toronto, Canada.

³ Department of Radiology, Children's Hospital, Vancouver, Canada.

⁴ Department of Radiology, Sick Children's Hospital, Toronto, Canada.

⁵ Department of Neurology, Val de Grace Hospital, Paris, France.

AJNR 8:185-192, March/April 1987.

© American Society of Neuroradiology.