

Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a *choice*.



[VIEW CATALOG](#)

AJNR

Erratum

AJNR Am J Neuroradiol 2025, 46 (4) 859

doi: <https://doi.org/10.3174/ajnr.A8711>

<http://www.ajnr.org/content/46/4/859>

This information is current as
of May 29, 2025.

In the article, “Low-Field Portable MR Imaging to Evaluate Ventricular Volumes: A Single-Center Retrospective Study” published in the August 2024 issue of *AJNR* (2024;45:1076–80; DOI: 10.3174/ajnr.A8269), Figs 1B and 1C were mislabeled. The correct image is shown below. The authors regret the error.




FIG 1. Comparison of SOC and portable clinical MR imaging as well as examples of manual segmentation in 1 participant, a 4-day-old boy without intracranial abnormalities such as ischemia or hemorrhage. Segmentation at 1 level of the lateral ventricles is shown here; SOC (A), SOC with segmentation (B), portable (C), portable with segmentation (D). Hyperfine scans were acquired with software, Version 8.1.0.1. TRA indicates transaxial; TSE, turbo spin echo; AXI, axial; R, right; L, left; P, posterior; A, anterior.

<http://dx.doi.org/10.3174/ajnr.A8711>