



Discover Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

Reply:

Y. Che

AJNR Am J Neuroradiol published online 19 November 2015

<http://www.ajnr.org/content/early/2015/11/19/ajnr.A4612.citation>

This information is current as of June 25, 2025.

REPLY:

Radiofrequency ablation (RFA) has been used for more than 20 years as a minimally invasive treatment of tumor. It has been widely recognized by scholars.^{1,2} In the past 10 years, RFA of thyroid nodules has developed rapidly because of the application of moving-shot technique, solving the problem of the important structures around the thyroid. A number of studies have shown that among the current treatment methods for benign thyroid nodules, RFA has many prominent advantages over the others, such as being minimally invasive, effective, relatively safe, cosmetically satisfactory, and having lower recurrence and so forth.^{3,4}

Obviously, however, how to select the RFA cases, not only to treat but also to identify them, was a key issue that concerned Bai et al. Based on a comprehensive analysis of the literature, the results of multicenter studies, and expert consensus, an RFA recommendation was published in 2012 by Korean Society of Thyroid Radiology (KSThR).⁵ In this publication, the indications for RFA of benign thyroid nodules included patients with nodule-related clinical problems: 1) symptoms of neck pain, dysphasia, foreign body sensation, discomfort, and cough; 2) cosmetic problems; and 3) autonomously functioning thyroid nodules causing problems related to thyrotoxicosis. Patients with nodules with a maximum diameter of >2 cm that continue to grow may be considered for thyroid RFA on the basis of symptoms and clinical concerns. The KSThR did not recommend thyroid RFA for follicular neoplasms or primary thyroid cancers because there is no evidence of treatment benefit. Before treatment, thyroid nodules should be confirmed as benign on at least 2 separate sonography-guided fine-needle aspirations and/or core needle biopsies.

At present, the concerning issue is the risk of malignancy in symptomatic nodular goiter. Ucler et al⁶ and Lee et al⁷ showed that the accuracy of fine-needle aspiration biopsy (FNAB) was 64.1%–99.6%. The diagnosis of false-negative findings was mainly due to groups of small cancer cells in the nodules and small cancers invisible under sonography. The results of the 2 punctures of the nodules in different places at different times should be benign, to avoid the risk of malignancy.⁸ Furthermore, with combined elastography or contrast-enhanced sonography, the puncture point and results are more accurate. The operation for benign thyroid nodules is thyroidectomy, and the identification standard is intraoperative frozen pathology. Prades et al⁹ reported that the accuracy of frozen pathology was 90% and maybe the potential malignancy was emerged in “benign” nodules. Negro et al¹⁰ reported that postoperative pathology of symptomatic nodular goiter accidentally confirmed microcarcinoma in 5%; papillary thyroid microcarcinoma (PTMC) was 96%, and there was the possibility of recurrence. With no difference from pathology, which determined the operation mode, FNAB was used for preoperative diagnosis in all minimally invasive treatments. Ito et al¹¹ reported that there was no obvious growth and metastasis in the 8-year follow-up without treatment of 732 cases of PTMC. Yue et al¹²

reported that during the 3-month follow-up period, ablation appears to be a safe and effective technique for solitary T1N0M0 PTMC.

Genetic testing had been used in the diagnosis of benign nodules undetermined by FNAB. Several molecular assays have been developed to detect the B-Raf proto-oncogene (*BRAF*) V600E mutation in fine-needle aspirates for the diagnosis of papillary thyroid cancer (PTC).¹³ Musholt et al¹⁴ considered that mutations of *RET/PTC*, *RAS*, and *PAX8*/peroxisome proliferator-activated receptor γ (*PPAR* γ) were predominantly associated with thyroid malignancy with varying frequency and had less impact on the clinical management. However, in the study of Song et al,¹⁵ *BRAF* mutations were the most common ones observed in PTCs, followed by *RET/PTC* rearrangements and *RAS* mutations, while follicular thyroid cancers were more likely to have *RAS* mutations or *PAX8/PPAR* γ rearrangements.¹⁵ Therefore, more extensive research is needed in genetic testing.

In conclusion, we suggested that RFA would be used as the first-line treatment of benign thyroid nodules with strict indications chosen. Elastography, contrast-enhanced sonography, and genetic testing would be used to differentiate the benign and malignant lesions.

REFERENCES

- Rossi S, Di Stasi M, Buscarini E, et al. Percutaneous radiofrequency interstitial thermal ablation in the treatment of small hepatocellular carcinoma. *Cancer J Sci Am* 1995;1:73–81 Medline
- Hefaidh R, Sabbah M, Ennaifer R, et al. Percutaneous treatment versus hepatic resection for the treatment of small hepatocellular carcinoma. *Tunis Med* 2014;92:711–16 Medline
- Deandrea M, Sung JY, Limone P, et al. Efficacy and safety of radiofrequency ablation versus observation for nonfunctioning benign thyroid nodules: a randomized controlled international collaborative trial. *Thyroid* 2015;25:890–96 CrossRef Medline
- Valcavi R, Tsamatopoulos P. Health-related quality of life after percutaneous radiofrequency ablation of cold, solid, benign thyroid nodules: a 2-year follow-up study in 40 patients. *Endocr Pract* 2015; 21:887–96 CrossRef Medline
- Na DG, Lee JH, Jung SL, et al; Korean Society of Thyroid Radiology (KSThR), Korean Society of Radiology. Radiofrequency ablation of benign thyroid nodules and recurrent thyroid cancers: consensus statement and recommendations. *Korean J Radiol* 2012;13:117–25 CrossRef Medline
- Ucler R, Usluogullari CA, Tam AA, et al. The diagnostic accuracy of ultrasound-guided fine-needle aspiration biopsy for thyroid nodules three centimeters or larger in size. *Diagn Cytopathol* 2015;43: 622–28 CrossRef Medline
- Lee YJ, Kim DW, Park YM, et al. Comparison of sonographic and cytological diagnoses of solid thyroid nodules: emphasis on the discordant cases. *Diagn Cytopathol* 2015 Sep 21. [Epub ahead of print] CrossRef Medline
- Singh Ospina N, Sebo TJ, Morris JC, et al. The value of repeat thyroid fine-needle aspiration biopsy in patients with a previously benign result: how often does it alter management? *Thyroid* 2015;25: 1121–26 CrossRef Medline
- Prades JM, Querat C, Dumollard JM, et al. Thyroid nodule surgery: predictive diagnostic value of fine-needle aspiration cytology and frozen section. *Eur Ann Otorhinolaryngol Head Neck Dis* 2013;130: 195–99 CrossRef Medline
- Negro R, Piana S, Ferrari M, et al. Assessing the risk of false-negative fine-needle aspiration cytology and of incidental cancer in nodular goiter. *Endocr Pract* 2013;19:444–50 CrossRef Medline
- Ito Y, Uruno T, Nakano K. An observation trial without surgical

- treatment in patients with papillary microcarcinoma of the thyroid. *Thyroid* 2003;13:381–87 [CrossRef Medline](#)
12. Yue W, Wang S, Yu S, et al. **Ultrasound-guided percutaneous microwave ablation of solitary T1N0M0 papillary thyroid microcarcinoma: initial experience.** *Int J Hyperthermia* 2014;30:150–57 [CrossRef Medline](#)
13. Choi R, Park KS, Kim JW, et al. **Evaluation of the Anyplex BRAF V600E real-time detection assay using dual-priming oligonucleotide technology in fine-needle aspirates of thyroid nodules.** *Ann Lab Med* 2015;35:624–29 [CrossRef Medline](#)
14. Musholt TJ, Musholt PB. **Molecular genetic markers for thyroid FNAB. Established assays and future perspective.** *Nuklearmedizin* 2015;54:94–100 [Medline](#)
15. Song YS, Lim JA, Park YJ. **Mutation profile of well-differentiated thyroid cancer in Asians.** *Endocrinol Metab (Seoul)* 2015;30:252–62 [CrossRef Medline](#)

Y. Che

Department of Ultrasound
First Affiliated Hospital of Dalian Medical University
Liaoning, China