



Providing Choice & Value
Generic CT and MRI Contrast Agents

**FRESENIUS
KABI**

CONTACT REP

AJNR

This information is current as
of July 29, 2025.

Spinal CSF Leaks: The Neuroradiologist Transforming Care

Mark D. Mamlouk, Andrew L. Callen, Ajay A. Madhavan, Niklas Lützen, Lalani Carlton Jones, Ian T. Mark, Waleed Brinjikji, John C. Benson, Jared T. Verdoorn, DK Kim, Timothy J. Amrhein, Linda Gray, William P. Dillon, Marcel M. Maya, Thien J. Huynh, Vinil N. Shah, Tomas Dobrocky, Eike I. Piechowiak, Joseph Levi Chazen, Michael D. Malinzak, Jessica L. Houk and Peter G. Kranz

AJNR Am J Neuroradiol published online 29 August 2024
<http://www.ajnr.org/content/early/2024/08/29/ajnr.A8484>

Spinal CSF Leaks: The Neuroradiologist Transforming Care

Mark D. Mamlouk, Andrew L. Callen, Ajay A. Madhavan, Niklas Lützen, Lalani Carlton Jones, Ian T. Mark, Waleed Brinjikji, John C. Benson, Jared T. Verdoorn, DK Kim, Timothy J. Amrhein, Linda Gray, William P. Dillon, Marcel M. Maya, Thien J. Huynh, Vinil N. Shah, Tomas Dobrocky, Eike I. Piechowiak, Joseph Levi Chazen, Michael D. Malinzak, Jessica L. Houk, Peter G. Kranz

ABSTRACT

SUMMARY: Spinal CSF leak care has considerably evolved over the past several years due to pivotal advances in its diagnosis and treatment. To the reader of the AJNR, it has been impossible to miss the exponential increase in groundbreaking research on spinal CSF leaks and spontaneous intracranial hypotension (SIH). While many clinical specialties have contributed to these successes, the neuroradiologist has been instrumental in driving this transformation due to innovations in non-invasive imaging, novel myelographic techniques, and image-guided therapies. In this editorial, we will delve into the exciting advancements in spinal CSF leak diagnosis and treatment and celebrate the vital role of the neuroradiologist at the forefront of this revolution, with particular attention to CSF leak related work published in the AJNR.

ABBREVIATIONS: SIH = spontaneous intracranial hypotension; CVF = CSF-venous fistula; CTM = CT myelography; DSM = digital subtraction myelography; CB-CTM = conebeam CT myelography; PCD-CT = photon counting detector CT

Received month day, year; accepted after revision month day, year.

From the Department of Radiology (M.D.M.), The Permanente Medical Group, Kaiser Permanente Medical Center, Santa Clara, Santa Clara, California; Department of Radiology and Biomedical Imaging (M.D.M., W.P.D. V.N.H.), University of California, San Francisco, San Francisco, California; Department of Radiology (A.L.C.), University of Colorado School of Medicine Anschutz Medical Campus, Aurora, Colorado; Department of Radiology (A.A.M., I.T.M., W.B., J.C.B., J.T.V., D.K.K.), Mayo Clinic, Rochester, Minnesota; Department of Neuroradiology (N.K.), University of Freiburg, Freiburg, Germany; Department of Radiology (L.C.J.), Guys and St Thomas's and Kings College Hospital Foundation Trusts, London, UK; Department of Radiology (T.J.A., L.G., M.D.M., J.L.H., P.G.K.) Duke University Hospital, Durham, North Carolina; Department of Imaging (M.M.M.), Cedars-Sinai Medical Center, Los Angeles, California; Department of Radiology (T.J.H.), Mayo Clinic, Jacksonville, Florida; Department of Diagnostic and Interventional Neuroradiology (T.D., E.I.P.), University of Bern, Inselspital, Bern, Switzerland; Department of Radiology (J.L.C.), Hospital for Special Surgery and Weill Cornell Medicine, New York, NY

The authors declare no conflicts of interest related to the content of this article.

Please address correspondence to Mark D. Mamlouk, MD, Department of Radiology, The Permanente Medical Group, Kaiser Permanente Medical Center, Santa Clara, 700 Lawrence Expy, Santa Clara, CA 95051; e-mail: mark.d.mamlouk@kp.org; @MarkMamloukMD

INTRODUCTION

Spinal CSF leak care has considerably evolved over the past several years due to pivotal advances in its diagnosis and treatment.¹ To the reader of the AJNR, it has been impossible to miss the exponential increase in groundbreaking research on spinal CSF leaks and spontaneous intracranial hypotension (SIH). While many clinical specialties have contributed to these successes, the neuroradiologist has been instrumental in driving this transformation due to innovations in non-invasive imaging, novel myelographic techniques, and image-guided therapies. In this editorial, we will delve into the exciting advancements in spinal CSF leak diagnosis and treatment and celebrate the vital role of the neuroradiologist at the forefront of this revolution, with particular attention to CSF leak related work published in the AJNR.

Spinal CSF Leak Types

Classifying the different types of spinal CSF leaks has contributed to our better understanding of the anatomy and pathophysiology and have led to advances in diagnosis and treatment. There are two main types of spontaneous spinal CSF leaks encountered in clinical practice: dural tears (ventral and lateral/posterior) and CSF-venous fistulas (CVFs).^{2, 3} Ruptured meningeal diverticula are an additional leak type that was discovered, but many diverticular leaks reported in the literature were likely lateral dural tears that mimicked a diverticular leak due to the arachnoid billowing through the dural defect.^{4, 5}

Spinal CSF Leak Publications

Spinal CSF leak research has been of recent topical interest in the neuroscience literature. To quantify this interest, we performed PubMed searches using the terms "spontaneous intracranial hypotension" or "CSF leak" or "CSF-venous fistulas" or "myelography" and "[journal name]" [journal]. The results were filtered between January 2020 and June 2024, and entries unrelated to spinal CSF leaks were excluded. Several neuroscience journals with known CSF leak publications were queried in the search, and any journal with 5 or more publications were included

(Table). *AJNR* had the highest number of publications (49), while *Headache* had the second highest with approximately one-third of the publications (16). Of the 11 total journals included, there were 5 radiology journals. The 49 *AJNR* spinal CSF leak publications constituted approximately 3% of all 1647 *AJNR* publications over the 3.5-year time frame. An analysis of spinal CSF leak publications in *AJNR* this millennium also shows a sharp increase in frequency over the past five years (Fig 1).

Not only are spinal CSF leak publications frequently published in the *AJNR*, but they are also commonly cited. On the *AJNR* homepage, under the recently “Most Cited” tab, 3 of the 5 articles listed are related to spinal CSF leaks,⁶⁻⁸ and there are 5 additional articles listed in the full *AJNR* list at the time of writing.⁹⁻¹⁴

Table: Number of Recent Spinal CSF Leak Publications by Medical Journal

Journal	Publications on CSF leak, CSF-venous fistula, or myelography related to SIH between January 2020-June 2024
<i>American Journal of Neuroradiology</i>	49
<i>Headache</i>	16
<i>Neurology</i>	13
<i>Clinical Neuroradiology</i>	12
<i>Interventional Neuroradiology</i>	12
<i>Neuroradiology</i>	10
<i>Journal of Neurointerventional Surgery</i>	9
<i>Cephalalgia</i>	7
<i>Neurology Clinical Practice</i>	7
<i>American Journal of Roentgenology</i>	6
<i>Journal of Neurosurgery Spine</i>	5
Others	<5

Note:--SIH indicates spontaneous intracranial hypotension




FIG 1. Number of Spinal CSF Leak Publications in *AJNR* from 2000-June 2024

Why All the Buzz on Spinal CSF Leaks?

One of the major reasons spinal CSF leak publications are topical is they can be challenging to diagnose and treat. Physicians of many specialties have struggles with this entity: how to make the diagnosis, how to localize a spinal leak, what treatment to pursue, and what to do when no leak is observed. There are 3 main research breakthroughs that have ameliorated some of these challenges and have propelled interest in the field. First, improved myelographic techniques have resulted in more precise localization of CSF leaks due to dural tears, allowing for more targeted and effective percutaneous and surgical treatments. Secondly, the discovery of CVFs has resulted in a paradigm shift in the approach to a patient with suspected SIH. CVFs represent an abnormal connection between the spinal subarachnoid space and an epidural vein, which abnormally shunts CSF into the venous system without an associated spinal epidural fluid collection. CVFs were first reported in 2014,¹⁵ but it was not until it was realized that lateral decubitus myelography resulted in a several fold increase in their detection that this diagnosis was made much more frequently in clinical practice.^{3, 6, 16, 17} Because of our improved understanding and better detection of CVFs, many patients in whom the CSF leak source could not be verified were subsequently diagnosed with a CVF.¹⁸ Though once thought to be rare, studies have suggested that CVFs are possibly the most common type of CSF leak in patients with SIH.¹⁰ The pathogenesis of CVFs is still being studied, but potential associations have been observed between preexisting intracranial hypertension as well as spinal degenerative disease.^{19, 20} Lastly, innovations in CVF treatment, driven by neuroradiologists,

further amplified excitement within the field of SIH. Surgery was the initial treatment modality described for CVFs, and while effective, is invasive and may require additional surgeries, as patients with CVFs may rarely develop same site recurrence or new CVFs after treatment. Subsequent work describing targeted fibrin glue occlusion and transvenous embolization were pioneered specifically for CVFs and resulted in significant enthusiasm given their minimally invasive nature.

Advanced Myelography

Prior to the myelographic work up, T2-weighted MR imaging of the spine is performed to evaluate the presence or absence of an extradural collection. This has largely replaced traditional myelography, whereby iodinated contrast is injected under fluoroscopy, allowed to diffuse evenly throughout subarachnoid space, and imaged after a substantial delay with the patient in the prone or supine position.²¹ While traditional myelography identifies the presence of an extradural fluid collection, it lacks the temporal resolution to precisely localize the site of dural defect. Instead, once a CSF leak is suspected, dynamic or decubitus CT myelography (CTM) or digital subtraction myelography (DSM) currently represent the two main modalities used for spinal CSF leak detection. If an extradural collection is observed on the spine MRI, a dural tear is suspected, and the patient is either positioned prone or decubitus Trendelenburg, dependent on the suspicion of leak type, typically using either a wedge or pillows. As an alternative, an innovative adjustable positioning device was invented to help achieve adequate angulation during myelography.²² After correct positioning, a lumbar puncture with contrast administration is performed, and rapid scanning is performed to capture the transition of contrast from the subarachnoid to extradural spaces, which represents the CSF leak site (Figs 2 and 3). CTM for dural tears is sometimes coined “ultrafast” or “dynamic” given that several successive temporally progressive acquisitions are performed while simultaneously injecting contrast to identify the leak site. More recently, single scan acquisitions with small contrast volumes have been described to achieve the same success with less radiation exposure.²³




FIG 2. Ventral dural tear on dynamic CT myelogram in a 34-year-old female showing the direct outflow of the contrast medium (open arrow in A, B) within seconds into the ventral epidural space at the T3-T4 level. The contrast medium then flows cranially within the epidural space (solid arrows in A). The underlying cause is a calcified disc at the T3-T4 level (open arrowhead in A, B).




FIG 3. Digital subtraction myelogram and conebeam CT of a lateral dural tear in a 36-year-old male. Digital subtraction myelogram in the right lateral decubitus position in anterior-posterior projection suggests a small epidural contrast medium egress, which is unchanged in the subsequent single radiograph (white solid arrows in A and B). C, Conebeam CT that follows a few minutes later confirms the epidural accumulation of contrast medium in coronal view (white solid arrows) at the right T12-L1 level. In addition, a cyst-like structure can be seen within the contrast collection, corresponding to an arachnoid layer (dashed black arrow in C) herniating through a lateral dural tear in the axilla of the exiting nerve root sleeve (black arrowhead in C), which was later confirmed by surgery.

Decubitus myelography with CTM or DSM currently serve as the mainstay imaging modalities for CVF detection after its initial discovery with prone technique. A CVF is suspected when the clinical and/or brain MRI features are suggestive of SIH but the spine MRI shows no extradural collection. When performing decubitus myelograms, contrast density and timing are complementary factors that help capture the egress of contrast from a distended spinal meningeal diverticulum into adjacent paraspinous veins.^{9, 24, 25} It was also described that maneuvers such as resisted inspiration while scanning may accentuate the CVF due to dynamic relationships between subarachnoid and venous pressures.^{8, 11, 26} CVFs have been described more frequently on the right side of the spine than the left, but there are no reliable ways to know which side the CVF will occur; therefore, evaluation of both sides may be needed. Right and left myelograms on separate days were reported initially,^{27, 28} while a same-day technique was later reported.¹² Additional tips that some authors have reported utility with for decubitus CTM are to perform more than one phase of scanning or to use real-time bolus tracking to improve detection.^{29, 30} Decubitus DSM and CTM both have unique advantages for CSF leak detection. While DSM has higher spatial and temporal resolution (Fig 4), CTM provides cross-sectional detail, has better contrast resolution, and is less susceptible to motion artifact in non-sedated patients. Conebeam CT myelography (CB-CTM) can be a helpful adjunct to DSM for CVFs or lateral dural tears.^{31, 32} The primary advantage of CB-CTM, when used in conjunction with DSM, is that it permits high-resolution cross-sectional imaging with a minimal delay between contrast injection and image acquisition. This can be helpful to clarify indeterminate findings on DSM, capitalizing on the advantages of both modalities (Fig 5). Dual energy CT and MRI with intrathecal gadolinium were also evaluated for CSF leaks.^{33, 34}




FIG 4. CSF-venous fistula on a digital subtraction myelogram (A) in a 47-year-old male with a right T5 CSF-venous fistula (arrow) that was treated with Onyx embolization (B). The pre-embolization brain MRI (C) demonstrated dural enhancement that nearly normalized one month after embolization (D).




FIG 5. Benefit of conebeam CT myelography for detection of a right T6 CSF-venous fistula. Select unsubtracted (A) and subtracted (B) images from a right lateral decubitus digital subtraction myelogram show faint linear paraspinal venous opacification (A, arrow). The finding is extremely difficult to appreciate on the unsubtracted image and essentially not seen on the subtracted image (B) due to a combination of pulmonary markings and respiratory motion. Axial (C) and coronal (D) maximum intensity projection images from a conebeam CT performed minutes later during active contrast injection show a clear right T6 CSF-venous fistula involving several lateral branch veins. In cases such as this, conebeam CT serves as an excellent adjunct to digital subtraction myelography.

Advances in Treatment

The greatest non-surgical innovations for CVF have been neuroradiologist-led treatments with fibrin occlusion and embolization, which were published in May 2021 and are now practiced throughout the world.

Historically, epidural patching for CVFs was not successful, but it was difficult to surmise adequate conclusions as to why, as many details were not clearly discussed, including the image guidance modality, the volume and type of injectate (fibrin glue or autologous blood), whether contrast was mixed with the injectate, and most importantly, was the patch performed along the CVF course or rather in the dorsal epidural space. CT-guided fibrin glue occlusion was successfully performed at a single center with effective results.^{35, 36} Subsequently, a multi-institutional and international study was performed, in which 59.7% of patients had complete clinical improvement and 34.5% had partial clinical improvement, with corresponding brain MRI improvement. One statistically significant conclusion from this study was that clinical improvement was observed if the injectate spread was concordant with the CVF drainage pattern (Fig 6).³⁷




FIG 6. CSF-venous fistula occlusion with targeted fibrin glue patching. A, Axial right decubitus CT myelogram shows a right T7 CSF-venous fistula with a paravertebral segmental vein (arrows). B, Axial CT treatment image demonstrates injected fibrin glue within the neural foramen and paravertebral space that matches the CSF-venous fistula drainage course, which is an important feature for treatment success. Post-treatment axial right decubitus CT myelogram (C) shows resolution of the CSF-venous fistula. The pretreatment brain MRI (D) demonstrated dural enhancement (arrows) that resolved one month after patching (E).

Spinal CVF embolization is a novel technique where the paraspinal vein is catheterized and a liquid embolic is injected resulting in a cast, thereby occluding the CSF-venous connection (Fig 4). It was first described at a single center with effective results clinically and on brain MRI and was subsequently published in larger studies.^{7, 38, 39} Besides the success with CVF embolization, the transvenous technique sparked greater interest and understanding of the paraspinal venous system, which was not well recognized. An atlas of the venous anatomy and guidance on where to approach the CVF throughout the spine has provided substantial guidance to the operating physician.⁴⁰

While both fibrin occlusion and embolization have markedly transformed spinal CSF leak treatment, surgery serves a vital role in patients with chronic dural tears and refractory CSF-venous fistulas and can be performed after a minimally invasive technique is attempted.^{4, 41}

The Promising Technologic Future

Photon counting detector CT (PCD-CT) will potentially serve an integral role in spinal CSF leak detection. Compared to traditional energy integrating detector (EID) CTs, PCD-CT demonstrates higher spatial and temporal resolution, along with inherent spectral sensitivity that can aid in detecting subtle CSF leaks without the need for dual energy/dual source techniques. In the context of spinal CSF leaks, CVFs were first evaluated by PCD-CT and showed high diagnostic yield.^{13, 42} One promising application is for CVFs that demonstrate drainage in the internal epidural venous plexus, an area that can be overlooked on myelography given challenges in spatial resolution.⁴³ High spatial resolution is also beneficial for detecting subtle venous opacification adjacent to meningeal diverticula and vertebral elements, where the juxtaposition of high-attenuation structures is difficult to resolve on EID CT (Fig 7). In addition to CVFs, PCD-CT has been shown to aid in the detection of CSF leaks from dural tears that also demands rapid scanning and high spatial resolution.⁴⁴ The high spatial resolution of PCD-CT can be maximally leveraged by using sharper quantitative kernels in image reconstruction, which can result in increased image noise. A deep learning algorithm has been applied to denoise PCD-CT images to enhance diagnostic quality, permitting the use of sharper kernels while retaining an acceptable signal-to-noise ratio (Fig 8).⁴⁵ As the technology becomes more widespread, PCD-CT will undoubtedly showcase additional benefits in CSF leak detection.




FIG 7. Advantage of high spatial resolution to detect a right T10 CSF-venous fistula on photon counting CT myelography. Axial and sagittal 0.2 mm images (A-B) from a right decubitus photon counting CT myelogram, reconstructed using a relatively smoother Br56 kernel, demonstrate a right T10 CSF-venous fistula involving the ventral and dorsal internal epidural venous plexus (A-B, arrows). Axial 0.2 mm images at the same level, reconstructed using both a smoother Br56 kernel (C) and a sharper Qr89 kernel with denoising (D) show involvement of the intervertebral vein that is only evident on the sharper Qr89 kernel. In some cases, maximizing spatial resolution using a sharper kernel with denoising is necessary to appreciate the full extent of venous opacification.




FIG 8. Use of denoised sharp kernel images on photon counting CT myelography for localization of a T5 ventral dural tear. Axial images from a prone dynamic photon counting CT myelogram, all at the same slice, timepoint, and window/level setting, demonstrate a ventral dural tear just to the right of midline (A-C, arrows). The precise location of the ventral leak is demonstrated with greater resolution when using a sharper Qr89 kernel (B-C) compared with a smoother Br56 kernel (A). Since the Qr89 kernel introduced noise into the image, a denoising algorithm was applied to permit the use of a sharper kernel while retaining an acceptable signal to noise ratio (C).

While there are many known clinical and imaging features of SIH, there are also many unknowns, including the underlying pathophysiologic state and when a spinal CSF leak will lead to symptoms and abnormal imaging findings. The brain MRI can occasionally be normal in the setting of a spinal CSF leak.⁴⁶ Studies on MR elastography and deep learning have shown promise in detecting underlying SIH in this context. In a pilot study, MR elastography found distinct stiffness and damping ratio patterns in SIH relative to controls.⁴⁷ While other brain MRI scoring systems evaluate the surface of the brain and downward morphologic sagging of the brainstem, MR elastography can evaluate changes in the brain parenchyma. In another study, an internally validated deep learning algorithm was created to identify patients with a CSF leak based on brain MR findings.⁴⁸ Additionally, glymphatic flow has been recently investigated in the setting of CSF-venous fistulas. A small case series demonstrated that patients with SIH may have impaired glymphatic clearance that could be restored after successful treatment.⁴⁹ These studies provide a glimpse into how advanced imaging techniques may provide increased sensitivity for the detection of SIH in patients who might have otherwise been inappropriately dismissed because of a conventionally normal brain MRI, as well as provide insight into the pathophysiologic underpinnings of spinal CSF leaks.

Redefining and Reaffirming Doctrines

The Monroe-Kellie doctrine posits that the combined volume of neuronal tissue, blood, and CSF is constant within the rigid skull, and any increase or decrease in one of these elements will lead to a reciprocal change in the others to maintain intracranial pressure homeostasis. However, studies have found that the calvarium in SIH may grow inward along the inner table of the skull resulting in layered hyperostosis, challenging the premise of this doctrine.^{50, 51} Conversely, the doctrine has been reaffirmed in the postsurgical setting in new ways such that in patients with surgical closure of a dural leak, ventricular CSF will increase with concomitant decrease in SIH brain findings.⁵²

After discovery of CVF, it became apparent they most commonly occur in the thoracic spine but rarely can occur in the lumbar and cervical spine. The revelation that sacral CSF-venous fistulas may also exist has added a new diagnostic area for evaluation.⁵³

Spontaneous intracranial hypotension can be a misnomer, as the opening pressure in many patients is within the normal range. In fact, pressures can even be elevated, particularly in patients with obesity.⁵⁴ This knowledge has diminished the role and need for opening pressure measurements for establishing a diagnosis of SIH.

Spinal and skull base CSF leaks are occasionally and erroneously categorized together, as it has been well established that spinal CSF leaks are associated with SIH and skull base leaks with intracranial hypertension. This distinction was recently reaffirmed in an analysis of skull base leaks while acknowledging that skull base leaks may rarely result in SIH.^{55, 56}

Redefining the Neuroradiologist

Spinal CSF leak care is a multidisciplinary effort. Various physician specialties, midlevel providers, nurses, technologists, and patient coordinators all contribute to the patient experience. While the neuroradiologist has always been intimately involved in this care team, the neuroradiologist's roles have propelled to the forefront. Because SIH is often first suggested on brain MRI, and because myelography is essential to precise spinal CSF leak diagnosis, it has been the neuroradiologist that has led efforts in optimizing the positioning techniques, scanning parameters, radiation safety, and new technologies to find the often-elusive spinal leak, which have transformed patients' lives and our profession. Due to the advances in patching and embolization, the neuroradiologist has served a more integral role as a treating physician rather than a diagnostic physician. Neuroradiologists have embraced these new roles with dedicated clinics, where in certain CSF leak centers the neuroradiologist is the primary physician contact for a CSF leak patient. While direct patient care is not new for neuroradiologists, these new avenues in CSF leak care have certainly expanded their broad armamentarium.

Patient-Centered Care

Diagnosis and treatment of spinal CSF leaks can occasionally be challenging. The entity is not well understood, is underdiagnosed, and has many clinical mimics.^{57, 58} Some patients are often dismissed and labeled incorrectly as having chronic pain or even psychiatric diagnoses, which can leave patients dejected and lose trust in the medical system. Because of this, many patients join support groups to seek advice from others with similar symptoms and to ultimately gain hope of recovery. Spinal CSF leak providers have recognized the patient role in this condition and have routinely embraced patient participation at their conferences where there have been physician and patient sessions to unite the two. Moreover, this converged at the inaugural "Bridging the Gap" conference in 2023 where patients shared their experiences, frustrations, and joys of their spinal leak journey.⁵⁹ These unique experiences have both strengthened the physician-patient relationship and, by providing a platform for patients to share their experiences, refined the diagnostic acumen of treating clinicians.

Our Job is Not Finished

While we are proud of these major developments in this spinal CSF leak era, there is much to be discovered and many questions are left unanswered. To date, research studies are retrospective and largely from individual institutions, which may be difficult to replicate at other centers with different resources. Prospective study designs, clinical trials, and comparative studies are needed to elevate spinal CSF leak care to new levels.⁶⁰ Furthermore, we need more specific clinical tests and scoring systems that are germane to the diagnosis to better triage which patients should undergo further workup to serve as an adjunct to existing imaging scoring tests.^{61, 62} Lastly, we need to increase awareness of this disease. Spinal CSF leak patients can enter the medical system through different venues and to various physicians, necessitating understanding of the main facets of this entity for prompt diagnosis and treatment.

Conclusions

Spinal CSF leak care hails many major accomplishments in diagnosis, treatment, and the patient experience. The neuroradiologist has played an integral role in its transformation, and journals like the *AJNR* serve as a beacon for disseminating these breakthroughs.

REFERENCES

1. Schievink WI. Spontaneous Intracranial Hypotension. *N Engl J Med*. 2021;385:2173-78
2. Schievink WI, Maya MM, Jean-Pierre S, Nuño M, Prasad RS, Moser FG. A classification system of spontaneous spinal CSF leaks. *Neurology*. 2016;87:673-9
3. Farb RI, Nicholson PJ, Peng PW, Massicotte EM, Lay C, Krings T, et al. Spontaneous Intracranial Hypotension: A Systematic Imaging Approach for CSF Leak Localization and Management Based on MRI and Digital Subtraction Myelography. *AJNR Am J Neuroradiol*. 2019;40:745-53
4. Schievink WI, Maya MM, Tay ASS, Taché RB, Prasad RS, Wadhwa V, et al. Lateral Spinal CSF Leaks in Patients with Spontaneous Intracranial Hypotension: Radiologic-Anatomic Study of Different Variants. *AJNR Am J Neuroradiol*. 2024
5. Madhavan AA, Farb RI, Brinjikji W, Cutsforth-Gregory JK, Schievink WI. Expounding on the Distinction between Lateral Dural Tears and Leaking Meningeal Diverticula in Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol*. 2024
6. Mamlouk MD, Ochi RP, Jun P, Shen PY. Decubitus CT Myelography for CSF-Venous Fistulas: A Procedural Approach. *AJNR Am J Neuroradiol*. 2021;42:32-36
7. Brinjikji W, Savastano LE, Atkinson JLD, Garza I, Farb R, Cutsforth-Gregory JK. A Novel Endovascular Therapy for CSF Hypotension Secondary to CSF-Venous Fistulas. *AJNR Am J Neuroradiol*. 2021
8. Amrhein TJ, Gray L, Malinzak MD, Kranz PG. Respiratory Phase Affects the Conspicuity of CSF-Venous Fistulas in Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol*. 2020
9. Mark I, Madhavan A, Oien M, Verdoorn J, Benson JC, Cutsforth-Gregory J, et al. Temporal Characteristics of CSF-Venous Fistulas on Digital Subtraction Myelography. *AJNR Am J Neuroradiol*. 2023;44:492-95
10. Mamlouk MD, Shen PY, Jun P, Sedrak MF. Spontaneous Spinal CSF Leaks Stratified by Age, Body Mass Index, and Spinal Level. *AJNR Am J Neuroradiol*. 2022;43:1068-72
11. Mark IT, Amans MR, Shah VN, Narsinh KH, Caton MT, Teixeira S, et al. Resisted Inspiration: A New Technique to Aid in the Detection of CSF-Venous Fistulas. *AJNR Am J Neuroradiol*. 2022;43:1544-47
12. Carlton Jones L, Goadsby PJ. Same-Day Bilateral Decubitus CT Myelography for Detecting CSF-Venous Fistulas in Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol*. 2022;43:645-48
13. Madhavan AA, Yu L, Brinjikji W, Cutsforth-Gregory JK, Schwartz FR, Mark IT, et al. Utility of Photon-Counting Detector CT Myelography for the Detection of CSF-Venous Fistulas. *AJNR Am J Neuroradiol*. 2023;44:740-44
14. AJNR Most Cited Articles <https://www.ajnr.org/articles/most-cited> Accessed June 27, 2024
15. Schievink WI, Moser FG, Maya MM. CSF-venous fistula in spontaneous intracranial hypotension. *Neurology*. 2014;83:472-3
16. Schievink WI, Maya MM, Moser FG, Prasad RS, Cruz RB, Nuño M, et al. Lateral decubitus digital subtraction myelography to identify spinal CSF-venous fistulas in spontaneous intracranial hypotension. *J Neurosurg Spine*. 2019;1-4
17. Kim DK, Brinjikji W, Morris PP, Diehn FE, Lehman VT, Liebo GB, et al. Lateral Decubitus Digital Subtraction Myelography: Tips, Tricks, and Pitfalls. *AJNR Am J Neuroradiol*. 2020;41:21-28
18. Behbahani S, Raseman J, Orlowski H, Sharma A, Eldaya R. Renal Excretion of Contrast on CT Myelography: A Specific Marker of CSF Leak. *AJNR Am J Neuroradiol*. 2020;41:351-56
19. Caton MT, Jr., Laguna B, Soderlund KA, Dillon WP, Shah VN. Spinal Compliance Curves: Preliminary Experience with a New Tool for Evaluating Suspected CSF Venous Fistulas on CT Myelography in Patients with Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol*. 2021;42:986-92
20. Sechrist EMZ, Pisani Petrucci SL, Andonov N, Lennarson P, Callen AL. The Spatial Relationship between Spinal Osteoarthritis and CSF Venous Fistulas in Patients with Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol*. 2024
21. Dobrocky T, Winklehner A, Breiding PS, Grunder L, Peschi G, Häni L, et al. Spine MRI in Spontaneous Intracranial Hypotension for CSF Leak Detection: Nonsuperiority of Intrathecal Gadolinium to Heavily T2-Weighted Fat-Saturated Sequences. *AJNR Am J Neuroradiol*. 2020;41:1309-15
22. Callen AL, Wojcik R, Bojanowski M. A Novel Patient-Positioning Device for Dynamic CT Myelography. *AJNR Am J Neuroradiol*. 2023;44:1352-55
23. Mamlouk MD, Shen PY, Dahlin BC. Modified Dynamic CT Myelography for Type 1 and 2 CSF Leaks: A Procedural Approach. *AJNR Am J Neuroradiol*. 2023;44:341-46
24. Gibby JT, Amrhein TJ, Young DS, Houk JL, Kranz PG. Diagnostic Yield of Decubitus CT Myelography for Detection of CSF-Venous Fistulas. *AJNR Am J Neuroradiol*. 2024
25. Mamlouk MD, Shen PY. Myelographic Timing Matters. *AJNR Am J Neuroradiol*. 2023;44:E16
26. Kranz PG, Malinzak MD, Gray L, Willhite J, Amrhein TJ. Resisted Inspiration Improves Visualization of CSF-Venous Fistulas in Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol*. 2023;44:994-98
27. Pope MC, Carr CM, Brinjikji W, Kim DK. Safety of Consecutive Bilateral Decubitus Digital Subtraction Myelography in Patients with Spontaneous Intracranial Hypotension and Occult CSF Leak. *AJNR Am J Neuroradiol*. 2020;41:1953-57
28. Shlapak DP, Kim DK, Diehn FE, Benson JC, Lehman VT, Liebo GB, et al. Time to Resolution of Inadvertent Subdural Contrast Injection during a Myelogram: When Can the Study Be Reattempted? *AJNR Am J Neuroradiol*. 2020;41:1958-62
29. Callen AL, Fakhri M, Timpone VM, Thaker AA, Dillon WP, Shah VN. Temporal Characteristics of CSF-Venous Fistulas on Dynamic Decubitus CT Myelography: A Retrospective Multi-Institution Cohort Study. *AJNR Am J Neuroradiol*. 2023;45:100-04
30. Huynh TJ, Parizadeh D, Ahmed AK, Gandia CT, Davison HC, Murray JV, et al. Lateral Decubitus Dynamic CT Myelography with Real-Time Bolus Tracking (dCTM-BT) for Evaluation of CSF-Venous Fistulas: Diagnostic Yield Stratified by Brain Imaging Findings. *AJNR Am J Neuroradiol*. 2023;45:105-12
31. Madhavan AA, Cutsforth-Gregory JK, Benson JC, Brinjikji W, Mark IT, Verdoorn JT. Conebeam CT as an Adjunct to Digital Subtraction Myelography for Detection of CSF-Venous Fistulas. *AJNR Am J Neuroradiol*. 2023;44:347-50
32. Lützen N, Demerath T, Volz F, Beck J, Urbach H. Conebeam CT as an Additional Tool in Digital Subtraction Myelography for the Detection of Spinal Lateral Dural Tears. *AJNR Am J Neuroradiol*. 2023;44:745-47

33. Huls SJ, Shlapak DP, Kim DK, Leng S, Carr CM. Utility of Dual-Energy CT to Improve Diagnosis of CSF Leaks on CT Myelography following Lateral Decubitus Digital Subtraction Myelography with Negative Findings. *AJNR Am J Neuroradiol.* 2022;43:1539-43
34. Chazen JL, Robbins MS, Strauss SB, Schweitzer AD, Greenfield JP. MR Myelography for the Detection of CSF-Venous Fistulas. *AJNR Am J Neuroradiol.* 2020;41:938-40
35. Mamlouk MD, Shen PY, Sedrak MF, Dillon WP. CT-guided Fibrin Glue Occlusion of Cerebrospinal Fluid-Venous Fistulas. *Radiology.* 2021:204231
36. Mamlouk MD, Shen PY, Dahlin BC. Headache response after CT-guided fibrin glue occlusion of CSF-venous fistulas. *Headache.* 2022;62:1007-18
37. Callen AL, Jones LC, Timponi VM, Pattee J, Scoffings DJ, Butteriss D, et al. Factors Predictive of Treatment Success in CT-Guided Fibrin Occlusion of CSF-Venous Fistulas: A Multicenter Retrospective Cross-Sectional Study. *AJNR Am J Neuroradiol.* 2023;44:1332-38
38. Brinjikji W, Garza I, Whealy M, Kissoon N, Atkinson JLD, Savastano L, et al. Clinical and imaging outcomes of cerebrospinal fluid-venous fistula embolization. *J Neurointerv Surg.* 2022
39. Brinjikji W, Madhavan A, Garza I, Whealy M, Kissoon N, Mark I, et al. Clinical and imaging outcomes of 100 patients with cerebrospinal fluid-venous fistulas treated by transvenous embolization. *J Neurointerv Surg.* 2023
40. Borg N, Cutsforth-Gregory J, Oushy S, Huynh T, Savastano LE, Cloft HJ, et al. Anatomy of Spinal Venous Drainage for the Neurointerventionalist: From Puncture Site to Intervertebral Foramen. *AJNR Am J Neuroradiol.* 2022
41. Schievink WI, Tache RB, Maya MM. Surgical Ligation of Spinal CSF-Venous Fistulas after Transvenous Embolization in Patients with Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol.* 2022;43:1073-76
42. Schwartz FR, Malinzak MD, Amrhein TJ. Photon-Counting Computed Tomography Scan of a Cerebrospinal Fluid Venous Fistula. *JAMA Neurol.* 2022;79:628-29
43. Madhavan AA, Cutsforth-Gregory JK, Brinjikji W, Bathla G, Benson JC, Diehn FE, et al. Diagnostic Performance of Decubitus Photon-Counting Detector CT Myelography for the Detection of CSF-Venous Fistulas. *AJNR Am J Neuroradiol.* 2023;44:1445-50
44. Madhavan AA, Cutsforth-Gregory JK, Brinjikji W, Benson JC, Johnson-Tesch BA, Liebo GB, et al. Benefits of Photon-Counting CT Myelography for Localization of Dural Tears in Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol.* 2024;45:668-71
45. Madhavan AA, Cutsforth-Gregory JK, Brinjikji W, Benson JC, Diehn FE, Mark IT, et al. Application of a Denoising High-Resolution Deep Convolutional Neural Network to Improve Conspicuity of CSF-Venous Fistulas on Photon-Counting CT Myelography. *AJNR Am J Neuroradiol.* 2023;45:96-99
46. Schievink WI, Maya MM, Tay ASS, Nisson PL, Acharya J, Taché RB, et al. Optic Nerve Sheath MR Imaging Measurements in Patients with Orthostatic Headaches and Normal Findings on Conventional Imaging Predict the Presence of an Underlying CSF-Venous Fistula. *AJNR Am J Neuroradiol.* 2024;45:655-61
47. Mark IT, Karki P, Cutsforth-Gregory J, Brinjikji W, Madhavan AA, Messina SA, et al. Evaluation of MR Elastography as a Noninvasive Diagnostic Test for Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol.* 2024;45:662-67
48. Faghani S, Moassefi M, Madhavan AA, Mark IT, Verdoorn JT, Erickson BJ, et al. Identifying Patients with CSF-Venous Fistula Using Brain MRI: A Deep Learning Approach. *AJNR Am J Neuroradiol.* 2024;45:439-43
49. Scharzt D, Finkelstein A, Zhong J, Brinjikji W, Bender MT. Improved Cerebral Glymphatic Flow after Transvenous Embolization of CSF-Venous Fistula. *AJNR Am J Neuroradiol.* 2024
50. Benson JC, Madhavan AA, Cutsforth-Gregory JK, Johnson DR, Carr CM. The Monroe-Kellie Doctrine: A Review and Call for Revision. *AJNR Am J Neuroradiol.* 2023;44:2-6
51. Babcock JC, Johnson DR, Benson JC, Kim DK, Luetmer PH, Shlapak DP, et al. Diffuse Calvarial Hyperostosis and Spontaneous Intracranial Hypotension: A Case-Control Study. *AJNR Am J Neuroradiol.* 2022;43:978-83
52. Dobrocky T, Rebsamen M, Rummel C, Häni L, Mordasini P, Raabe A, et al. Monroe-Kellie Hypothesis: Increase of Ventricular CSF Volume after Surgical Closure of a Spinal Dural Leak in Patients with Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol.* 2020;41:2055-61
53. Mark IT, Morris PP, Brinjikji W, Madhavan AA, Cutsforth-Gregory JK, Verdoorn JT. Sacral CSF-Venous Fistulas and Potential Imaging Techniques. *AJNR Am J Neuroradiol.* 2022;43:1824-26
54. Schievink WI, Maya M, Prasad RS, Wadhwa VS, Cruz RB, Moser FG. Spinal CSF-Venous Fistulas in Morbidly and Super Obese Patients with Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol.* 2021;42:397-401
55. Mark IT, Cutsforth-Gregory J, Luetmer PH, Madhavan AA, Oien MP, Farnsworth PJ, et al. Skull Base CSF Leaks: Potential Underlying Pathophysiology and Evaluation of Brain MR Imaging Findings Associated with Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol.* 2024
56. Schievink WI, Michael LM, 2nd, Maya M, Klimo P, Jr., Elijovich L. Spontaneous Intracranial Hypotension Due to Skull-Base Cerebrospinal Fluid Leak. *Ann Neurol.* 2021;90:514-16
57. Bond KM, Benson JC, Cutsforth-Gregory JK, Kim DK, Diehn FE, Carr CM. Spontaneous Intracranial Hypotension: Atypical Radiologic Appearances, Imaging Mimickers, and Clinical Look-Alikes. *AJNR Am J Neuroradiol.* 2020;41:1339-47
58. Benson JC, Madhavan AA, Mark IT, Cutsforth-Gregory JK, Brinjikji W, Verdoorn JT. Likelihood of Discovering a CSF Leak Based on Intracranial MRI Findings in Patients without a Spinal Longitudinal Extradural Collection: A New Probabilistic Scoring System. *AJNR Am J Neuroradiol.* 2023;44:1339-44
59. Callen AL, Petrucci Pisani SL, Lennarson P, Birlea M, MacKenzie J, Buchanan AJ. Perspectives from the Inaugural "Spinal CSF Leak: Bridging the Gap" Conference: A Convergence of Clinical and Patient Expertise. *AJNR Am J Neuroradiol.* 2024
60. Amrhein TJ, Williams JW, Jr., Gray L, Malinzak MD, Cantrell S, Deline CR, et al. Efficacy of Epidural Blood Patching or Surgery in Spontaneous Intracranial Hypotension: A Systematic Review and Evidence Map. *AJNR Am J Neuroradiol.* 2023;44:730-39
61. Houk JL, Morrison S, Peskoe S, Amrhein TJ, Kranz PG. Validity of the Bern Score as a Surrogate Marker of Clinical Severity in Patients with Spontaneous Intracranial Hypotension. *AJNR Am J Neuroradiol.* 2023;44:1096-100