

Table 2: Visualization of specific arterial segments, noise, and artifacts, PCD-CT vs. EID-CT.

	Cavernous ICA (-2 to 2)	Ophthalmic (-2 to 2)	M3 (-2 to 2)	A1 (-2 to 2)	A3 (-2 to 2)	P1 (-2 to 2)	P3 (-2 to 2)	Noise (-2 to 2)	Artifacts (-2 to 2)
Session 1: PCD-SR vs EID-SR									
R1	1±0.2	1.2±0.4	1±0	1±0.3	1±0	1±0.2	1.3±0.5	1±0	1±0
R2	1±0	1±0	1.8±0.4	1±0	1.3±0.4	1±0	1.4±0.5	1.6±0.5	0.9±0.3
Combined	1±0.1	1.1±0.3	1.4±0.5	1±0.2	1.1±0.3	1±0.1	1.3±0.5	1.3±0.5	1±0.2
Session 2: PCD-HR-CNN vs EID-SR									
R1	1±0	1.4±0.5	1±0.2	1±0.2	1±0	0.9±0.3	1.2±0.4	1±0	1±0
R2	1.9±0.4	2±0	2±0	2±0.2	2±0	2±0.2	2±0.2	1.9±0.3	-0.4±0.8
Combined	1.4±0.5	1.7±0.5	1.5±0.5	1.5±0.5	1.5±0.5	1.4±0.6	1.6±0.5	1.4±0.5	0.3±0.9