

Online Supplemental Data: Bivariate analysis comparing patients with incidental brain aneurysms on screening MRA with those who didn't

Characteristics	Full Sample (N=24397)	No Incidental Aneurysm (N=22313)	Incidental Aneurysm (N=2084)	P Value
Sex, N (%)				<0.001
Female	14862 (60.9%)	13463 (60.3%)	1399 (67.1%)	
Male	9535 (39.1%)	8850 (39.7%)	685 (32.9%)	
Age				<0.001
Mean ± SD	60.4 ± 17.4	60 ± 17.6	64 ± 15.4	
Median (IQR)	62 (49, 73)	61 (49, 73)	65 (54, 76)	
Race & Ethnicity, N (%)				0.045
Non-Hispanic White	2953 (12.1%)	2727 (12.2%)	226 (10.8%)	0.071
Non-Hispanic Black	8194 (33.6%)	7503 (33.6%)	691 (33.2%)	0.683
Hispanic	8758 (35.9%)	7953 (35.6%)	805 (38.6%)	0.007
Other	3798 (15.6%)	3497 (15.7%)	301 (14.4%)	0.147
Unknown	694 (2.8%)	633 (2.8%)	61 (2.9%)	0.867
Constipation, N (%)	3946 (16.2%)	3572 (16.0%)	374 (17.9%)	0.023
Renal Disease, N (%)	5508 (22.6%)	5028 (22.5%)	480 (23.0%)	0.622
ADPCKD, N (%)	60 (0.2%)	55 (0.2%)	5 (0.2%)	1.000
HTN, N (%)	16336 (67.0%)	14833 (66.5%)	1503 (72.1%)	<0.001
Hyperlipidemia, N (%)	11162 (45.8%)	10123 (45.4%)	1039 (49.9%)	<0.001
Cocaine Use, N (%)	431 (1.8%)	397 (1.8%)	34 (1.6%)	0.687

Alcohol Use, N (%)	792 (3.2%)	724 (3.2%)	68 (3.3%)	1.000
Smoking History, N (%)	5058 (20.7%)	4594 (20.6%)	464 (22.3%)	0.076

ADPCKD – Autosomal Dominant Polycystic Kidney Disease, HTN – Hypertension, IQR – Interquartile Range, SD – Standard Deviation